

ON THE SUPERCONDUCTING PHASES OF UTe_2


V.I. Marchenko^a, L.A. Melnikovsky^{b,a*}

^a P.L. Kapitza Institute for Physical Problems, Moscow, Russia

^b Weizmann Institute of Science, Rehovot, Israel

The anomalous triple point discovered in superconducting UTe_2 may indicate a gauge invariance breaking in the high-pressure phase, manifesting as Bose condensation of bound states of eight or even more electrons.

On the P, T diagram of UTe_2 [1], three lines of continuous phase transitions between the normal phase n and two distinct superconducting phases, s_1 and s_2 , meet at a single point (Fig.a). To our knowledge, such a point has not been observed in other systems, although a tetracritical one (Fig.b), as predicted by Landau in 1937 does exist. A satisfactory explanation for this discovery is still lacking [2]. In this note, we point out that a natural extension of the concept of gauge invariance breaking may facilitate such an explanation.



It was recently found [3] that the magnetic flux quantum in superconducting CsV_3Sb_5 changes with increasing temperature, taking the values $h/(2ek)$, where $k = 1, 2, 3$. This has been interpreted as evidence for the formation of $2k$ -electron condensates.

Let us consider, within Landau theory of continuous transitions, a tetracritical point (Fig.b) in a superconductor. Assuming zero quasimomentum condensates, we restrict our consideration to its crystal class D_{2h} . Let ψ correspond to a $2k$ -electron condensate and η to a $2mk$ -electron condensate, where $m > 1$. It is necessary to add to the usual Ginzburg-Landau terms:

$$(at + vp)|\psi|^2 + A|\psi|^4 + (bt - wp)|\eta|^2 + B|\eta|^4, \quad (1)$$

where p, t are deviations of pressure and temperature from the intersection, biquadratic and Indenbom-

type [4] crossterms:

$$C|\psi|^2|\eta|^2 + I(\psi^{*m}\eta + \psi^m\eta^*). \quad (2)$$

The latter is allowed if η transforms according to the unit representation of the crystal class for even m , or according to the same representation as ψ for odd m . In this case, η is induced by ψ as an improper parameter [4], and the transition line $n|\eta$ terminates at the intersection point, which thus becomes a triple point. For $m = 2$, however, the Indenbom's term is cubic in ψ, η at the intersection itself, and superconductivity emerges discontinuously at higher temperature. In the case $m = 3$, it is also cubic in ψ on the line $\eta|\psi\eta$. Only for $m > 3$ does this scenario agree with the anomalous phase diagram of UTe_2 . Note that Indenbom's invariant, though small here, is linear in η and acts as a conjugate field, thus leading to the disappearance of one branch from the phase diagram. The biquadratic term produces a kink between the $n|s_1$ and $s_2|s_1$ lines, but the available experimental accuracy is insufficient to identify even its sign.

We thank M. Elbaum, I.A. Fomin, E.I. Katz, and E.V. Surovtsev for helpful discussions.

1. D. Braithwaite, M. Vališka, G. Knebel, et al, Communications Physics 2, 147 (2019)
2. T. Vasina, D. Aoki, A. Miyake, et al, Phys. Rev. Lett. 134, 096501 (2025)
3. J. Ge, P. Wang, Y. Xing, et al, Phys. Rev. X14, 021025 (2024)
4. L.D. Landau, E.M. Lifshitz, Electrodynamics of Continuous Media, §20, Pergamon Press, Oxford (1984)

* E-mail: lev.melnikovsky@weizmann.ac.il